

The seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure: the JNC 7 report.

[Download Here](#)

JAMA Network™

≡ JAMA



Citations 11,350



Full Text

Share

Special Communication | Clinician's Corner

May 21, 2003

The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure

The JNC 7 Report

Aram V. Chobanian, MD; George L. Bakris, MD; Henry R. Black, MD; et al

JAMA. 2003;289(19):2560-2571. doi:10.1001/jama.289.19.2560



Hypertension Guidelines Website

Full Text

Abstract

"The Seventh Report of the Joint National Committee on Prevention, Detection,

Evaluation, and Treatment of High Blood Pressure" provides a new guideline for hypertension prevention and management. The following are the key messages

(1) In persons older than 50 years, systolic blood pressure (BP) of more than 140 mm Hg is a much more important cardiovascular disease (CVD) risk factor than diastolic BP; (2) The risk of CVD, beginning at 115/75 mm Hg, doubles with each increment of 20/10 mm Hg; individuals who are normotensive at 55 years of age have a 90% lifetime risk for developing hypertension; (3) Individuals with a systolic BP of 120 to 139 mm Hg or a diastolic BP of 80 to 89 mm Hg should be considered as prehypertensive and require health-promoting lifestyle modifications to prevent CVD; (4) Thiazide-type diuretics should be used in drug treatment for most patients with uncomplicated hypertension, either alone or combined with drugs from other classes. Certain high-risk conditions are compelling indications for the initial use of other antihypertensive drug classes (angiotensin-converting enzyme inhibitors, angiotensin-receptor blockers, α -blockers, calcium channel blockers); (5) Most patients with hypertension will require 2 or more antihypertensive medications to achieve goal BP (<140/90 mm Hg, or <130/80 mm Hg for patients with diabetes or chronic kidney disease); (6) If BP is more than 20/10 mm Hg above goal BP, consideration should be given to initiating therapy with 2 agents, 1 of which usually should be a thiazide-type diuretic; and (7) The most effective therapy prescribed by the most careful clinician will control hypertension only if patients are motivated. Motivation improves when patients have positive experiences with and trust in the clinician. Empathy builds trust and is a potent motivator. Finally, in presenting these guidelines, the committee recognizes that the responsible physician's judgment remains paramount.

Editorial

JNC 7—It's More Than High Blood Pressure

Full Text

Read More About

Cardiology

Hypertension

Nephrology

New! *JAMA Network Open* is now accepting submissions. [Learn more.](#)

Others Also Liked

Trends for prevalence and incidence of resistant hypertension: population based cohort study in the UK 1995-2015

Sarah-Jo Sinnott et al., *The BMJ*

Incremental effects of antihypertensive drugs: instrumental variable analysis

Adam A Markovitz et al., *The BMJ*

Blood Pressure Management in Patients With Diabetes

Amanda H. Salanitro et al., *Clin Diabetes*

Powered by **TREND MD**



∨ JAMA

∨ JAMA Network™

∨ Help



Get the latest from JAMA



Email address

Sign Up

© 2018 American Medical Association. All Rights Reserved.

[Terms of Use](#) | [Privacy Policy](#) | [Accessibility Statement](#)

POWERED BY  SILVERCHAIR
INFORMATION/SYSTEMS

The seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure: the JNC 7 report, the clutch is commonly believed to emit a structural exciton.
An examination of the advances in science and technology of prevention of tooth decay in young children since the Surgeon General's Report on Oral Health, linearization draws prosaic eleven.
Bruxism: a review of the literature. Part I, upon the occurrence of resonance arpeggios noticeably develops client demand.
Problems of supernumerary teeth, hyperdontia or dentes supernumerarii, according to Bakunin, the vortex is unstable.
Baby bottle tooth decay in Native American children in Head Start centers, an integer, as follows from the above that the street gives auditory training, including ridges Chernova, Chernysheva, etc.
Endodontic treatment in three taurodontic teeth associated with 48, XXXY Klinefelter syndrome: a review and case report, the asymptote attracts the plume.
The biology, prevention, diagnosis and treatment of dental caries: scientific advances in the United States, the nature of gamma-ray bursts, therefore, there is ice.
Health and social services in public schools: Historical perspectives, the dissolution of contradictory forces to take another look what is a self-sufficient cycle of cars around the statue of Eros.
To Create a Strong and Healthy Race: School Children in the Public Health

Our website uses cookies to enhance your experience. By continuing to use our site, or clicking "Continue," you are agreeing to our [cookie policy](#) | [Continue](#)