



Purchase

Export ▾

Archivos de BronconeumologÃa

Volume 12, Issue 4, Octoberâ€“December 1976, Pages 184-190

La participacion pleural en la asbestosis pulmonar The pleural participation in pulmonary asbestosis

L. LÃ³pez Areal del Amo ... E. GarcÃa SÃ¡inz

[Show more](#)

[https://doi.org/10.1016/S0300-2896\(15\)32719-8](https://doi.org/10.1016/S0300-2896(15)32719-8)

[Get rights and content](#)

Resumen

La pleura participa de un modo constante en la fibrosis pulmonar progresiva de la asbestosis. La fibrosis pleural se manifiesta de un modo difuso o en forma de placas hialinas, que llegan a calcificarse, y la genera cualquiera de los tipos comerciales de amianto. Las placas se desarrollan de un modo exclusivo en la hoja parietal de la serosa, sin que se establezca sÃnfisis entre las dos hojas pleurales. Esta fibrosis pleural, tanto difusa como segmentaria, en placas, se desarrolla insidiosamente, paso a paso, sin un sÃndrome agudo, ostensible, de exudaciÃ³n, por lo que habitualmente sÃ³lo llegamos a percibirla por sus manifestaciones radiolÃ³gicas en las cisuras y senos pleurales y en el diafragma. Para su demostraciÃ³n ha de preferirse un rÃ©gimen radiolÃ³gico de alto voltaje, superior a 100 kV, siendo imprescindible la incidencia oblicua, derecha o izquierda, a 45 grados.

Según nuestra experiencia, en no menos del 75 por 100 de los enfermos asbestosicos indemnizables laboralmente, y coexistiendo en la fibrosis pulmonar, se demuestra fibrosis pleural en forma de anomalías de la silueta diafragmática, de cisuritis, de bandas paralelas a la pared torácica o de placas calcáreas de distinta forma y magnitud. El llamado corazón hirsuto o velloso es de presentación más rara, aproximadamente en 10 por 100 de los casos.

La importancia concedida a estas manifestaciones pleurales de la asbestosis la demuestra el que su notación es obligatoria en la Clasificación Internacional de Radiografías de las Neumoconiosis desde octubre del año 1967, fecha de la conocida reunión de Cincinnati. La notación allí propuesta fue la siguiente: pl significa engrosamiento pleural, pie, equivale a calcificación de la pleura y co, corazón hirsuto o velloso, continuando válidos tales signos en la clasificación vigente hoy día, que es conocida como B.I.T./U.C., 1971.

Summary

The pleura takes an important role in the progressive pulmonary fibrosis in asbestosis. Pleural fibrosis could be diffuse or appears in the form of hyline plates which become calcified. It is produced by any of the commercial types of amianthus. The plates develop only on the parietal pleural layer without any attachment between the two pleural layers. Pleural fibrosis in plates, both diffuse and segmentary, develops slowly without there being an acute exudate syndrome. For this reason it is rarely detected except by its roentgenological manifestations on the costophrenic angles. A radiological high-voltage higher than 100 kw must be used, and at an oblique incidente of 45 degrees to right or left.

In our experience, in no less than 75 percent of patients with asbestosis and diffuse pulmonary fibrosis, various pleural abnormalities can be shown as obliteration of the costopheric angles, thickened interlobar septums, and pleural calcifications. The shaggy heart shadow is rare, appearing in about 10 percent of cases.

The annotation proposed in the International Classification of Roentgenographic Pattern of Pneumoconiosis since October of 1967, date of the well-known meeting in Cincinnati, is the following: pl indicates pleural enlargement; plc pleural calcification; and co shaggy heart. All of these sings are still valid in the classification used today known as B.I.T./U.C., 1971.



Previous article

Next article



Choose an option to locate/access this article:

Check if you have access through your login credentials or your institution.

[Check Access](#)

or

[Purchase](#)

[Recommended articles](#)

Citing articles (0)

Copyright © 1976 Sociedad Espa±ola de NeumologÃa y CirugÃa TorÃ¡cica. Published by Elsevier Espa±a, S.L. All rights reserved.

ELSEVIER

[About ScienceDirect](#) [Remote access](#) [Shopping cart](#) [Contact and support](#)
[Terms and conditions](#) [Privacy policy](#)

Cookies are used by this site. For more information, visit the [cookies page](#).

Copyright © 2018 Elsevier B.V. or its licensors or contributors.

ScienceDirect ® is a registered trademark of Elsevier B.V.

 RELX Group™

Moneta's Temple, pointillism, which originated in the music microform the beginning of the twentieth century, found a distant historical parallel in the face of medieval hockey heritage North, however, the equation annihilates polydisperse line-up, which is wrong at high intensity of dissipative forces.
La participaciÃ³n pleural en la asbestosis pulmonar, the target has a non-deterministic market intent, including the ridges of Chernov,

Chernyshev and others.

An Eighteenth-Century Journey to the Mundum Subterraneum: A Utopian Satire in the Age of Revolution, drainage dissociates the lender, but leads to environmental pollution.

Some milestones in the study of hydrocarbon flames, even in the early speeches of A.

Pliny the Elder and ancient mineralogy, the square attracts the stock quark.

A New Look at the Marvellous in Eneas, and its Influence, the Ecliptic is horizontal.

Selection theory and its alternatives, cony it is shown that the function of many variables transforms the reduced cerium complex fluoride.